



Pharmacology



Drug Fundamentals, Plus the Most Frequently Prescribed Drug Classifications—including Indications, Reactions, Examples & More

definitions

pharmacodynamics Study of the mechanisms of action of drugs within the body and how drugs produce their effects in the body

pharmacogenetics Study of drug reactions in the body that are unanticipated or unusual, and may have a hereditary basis for the response

pharmacokinetics Study of drug actions as they move through the body; the way the body absorbs, distributes, metabolizes and excretes drugs; mathematical study of drugs based on time and dose

pharmacology Study of biologically active compounds, how they react in the body and how the body reacts to them


pharmacotherapeutics Study of drugs used to prevent, treat or diagnose disease

pharmacy Preparation and dispensing of drugs

toxicology Study of harmful or poisonous effects of drugs

drug names

DRUG



pharmacodynamics

Receptors

- Protein molecules with one or more binding sites, located on cell membranes
- Receive a signal from the body's chemicals: neurotransmitters, hormones, enzymes
- Signal will cause a molecular event on the inside of the cell to occur
- Drugs** Enhance (*agonist*), diminish (*partial agonist*) or block (*antagonist*) the generation, transmission or receiving of the signal
- Affinity** Attraction between a drug and a receptor
- High affinity** Drug will bind easily to the receptor
- Low affinity** Requires a higher concentration of the drug to get a therapeutic response

Drug Potency

- Amount of drug required to produce a therapeutic response

Dose Response Curve

- Effective Dose (ED)** Amount of drug that produces a therapeutic response in 50% of the people taking it
- Toxic Dose (TD)** Amount of drug that produces adverse effects in 50% of the people taking it
- Therapeutic Index (TI)** Margin of safety; ratio between the TD and the ED
- The higher the TI**, the safer the drug is considered to be; in general, nonprescription drugs have much higher TIs than prescription drugs

pharmacokinetics

Routes drugs take to get into the body

- Enteral:**
 - Enters the body through the GI tract
 - Taken by mouth, through the rectum, under the tongue or held in the cheeks
- Parenteral:**
 - Enters the body through a different means (i.e., other than the GI tract)
 - Can be injected into the veins, arteries, muscles, spinal cord, or under the skin; inhaled through the lungs; transdermally through the skin via ointment or patch

Absorption

- Bioavailability** Percent absorbed into systemic circulation after administration
 - Bioavailability depends on route of administration as well as the drug's ability to cross membranes and reach its target
 - First Pass Effect:**
 - Drugs absorbed through the stomach and small intestine must pass through the liver before circulating systemically
 - Liver can inactivate the drug, making less of the drug available to reach the target organ
- Absorption** at cellular level occurs through passive transport, active transport, pinocytosis and facilitated diffusion

Distribution

- Influenced by several factors:**
 - Tissue permeability:** Ability of drug to pass through the membranes rapidly affects the extent to which the drug moves around in the body
 - Blood flow:** Once in the blood stream, will get to the organs and tissues that are highly perfused
 - Plasma proteins binding:** Drug can bind to a protein that will render the drug inactive; only an unbound drug can attach to the receptors
 - Binding to subcellular components
 - blood pH

Drug Storage Sites

- Adipose tissue** Primary site; lipid-soluble; drugs tend to remain for long periods of time due to low metabolic rates of drugs and poor blood perfusion of tissue
- Bone** Toxic agents like heavy metals
- Muscle** Binding can cause muscle to store drugs
- Organs** Liver and kidneys

Metabolism

- Biotransformation** Chemical changes that occur in the drug following administration
- Metabolite** Altered version of the chemical compound
- Can have a higher or lower rate of activity than the original drug; if higher, drug is given as an inactive or prodrug form
- Prodrug** Requires metabolism or activation of drug in order for it to act within the body

Excretion

- First order** Rate of removal of drug from the body is proportional to the concentration of the drug in the plasma
- Half-life** Time required to decrease the blood levels of a drug by one-half
- A one-time drug dosage will be eliminated almost completely by 5 half-lives
- A drug given on a continuous dosage schedule will reach steady state concentration after 5 half-lives
- Steady state** Rate of drug administration is equal to the rate of drug excretion
- Organs that excrete drugs** Kidneys, lungs, sweat glands, mammary glands, salivary glands, skin and GI tract

schedule drugs



Schedule Class

Schedule 1: C-1
Schedule 2: C-II

Schedule 3: C-III

Schedule 4: C-IV

Schedule 5: C-V

Characteristics

- High abuse potential; not legal; no acceptable medical use; no prescriptions available
- High abuse potential and severe dependence liability; current, accepted medical use; prescription drug-signed; not stamped prescription; 30-day supply, no refills
- Less abuse potential; low-moderate physical dependence; high psychological dependence; by prescription only, expires within 6 months; max. 5 refills on one script
- Less abuse potential than C-III drugs; accepted medical use; limited physical and psychological dependence; written or verbal prescription, expires in 6 months; max. 5 refills on one script
- Limited abuse potential; accepted medical use; small amounts of narcotics used as antitussives (cough medicine) or antidiarrheals; may not need a prescription but must be recorded as a transaction

Examples (C-I to C-IV)

- Heroin, LSD, cocaine, marijuana, methaqualone
- Opium, morphine, coca, methadone
- Amphetamines, codeine, barbiturates, Valium, Xanax, anabolic steroids
- Chloral hydrate, meprobamate, paraldehyde, phenobarbital

Pharmaceutical Classifications

adrenergics

Mimic naturally occurring catecholamines (epinephrine, norepinephrine and dopamine) or stimulate the release of norepinephrine

Indications Alpha-adrenergic agonists used to treat hypotension

⊙ **Common drug examples:**

- **Norepinephrine** Lovophed
- **Pseudoephedrine** Cenafed, Dimetapp, Sudafed, Triaminic DM (OTC used to treat other conditions)

⊙ **Adverse reactions:** Increased blood pressure, AV block; other effects include: nausea, vomiting, sweating, goose bumps, rebound miosis, difficulty in urinating, headache, dilated pupils, photophobia, burning, stinging and blurry eyes

Beta 1 adrenergic agonists Bradycardia, low cardiac output, paroxysmal atrial or nodal tachycardia, ventricular fibrillation, cardiac output

⊙ **Common drug examples:**

- **Dobutamine hydrochloride** Dobutrex

⊙ **Adverse reactions** Tachycardia, palpitations and other arrhythmias, premature and ventricular contractions, tachyarrhythmias and myocardial necrosis

Beta 2 adrenergic agonists Acute and chronic bronchial asthma, emphysema, bronchitis, acute hypersensitive (allergic) reaction to drugs, delays delivery in premature labor, dysmenorrhea

⊙ **Common drug examples:**

- **Albuterol sulfate** Proventil, Ventolin, Volmax
- **Bitolterol mesylate** Tornalate
- **Metaproterenol sulfate** Alupent
- **Pirbuterol acetate** Maxair
- **Salmeterol xinafoate** Serevent
- **Terbutaline** Brethine, Bricanyl

⊙ **Adverse reactions** Nervousness, tremors, headaches, tachycardia, palpitations, hypertension, nausea, vomiting, cough

Dopamine Improves blood flow to the kidneys; used in acute renal failure, heart failure and shock

⊙ **Common drug examples:**

- **Dopamine hydrochloride** Intropin

⊙ **Adverse reactions** Headaches, ectopic beats, tachycardia, hypotension, bradycardia, nausea, vomiting, hyperglycemia, asthma attacks, anaphylactic reactions

adrenocorticoids

Glucocorticoids Regulate carbohydrate, lipid and protein metabolism; block inflammation; regulate body's immune response

⊙ **Indications** Asthma, advance pulmonary tuberculosis, pericarditis, acute and chronic inflammation, adrenal insufficiency, antenatal use in preterm labor, hypercalcemia, cerebral edema, acute SCI, MS, shock

⊙ **Common drug examples:**

- **Betamethasone** Beclvent, QVAR, Vanceryl
- **Hydrocortisone** Cortet, Hycort
- **Methylprednisone** Medrol, Meprolone, Metacort
- **Prednisone** Apo-prednisone, Deltasone, Meticort, Orasone, Sterapred
- **Triamcinolone** Azmacort, Nasacort

⊙ **Adverse reactions** Primarily a catabolic effect on muscle, bone, ligament, tendon; suppression of hypothalamic-pituitary-adrenal pathway; Cushingoid syndrome with long-term use; other effects include euphoria, insomnia, psychotic behavior, pseudotumor, mental changes, nervousness, restlessness, heart failure, hypertension, edema, acute tendon ruptures, delayed wound healing

- **Withdrawal symptoms if drugs stopped abruptly** Fever, myalgias, arthralgias, malaise, nausea, orthostatic hypotension, dizziness, fainting, dyspnea, hypoglycemia

Mineralocorticoid Regulates electrolyte homeostasis

⊙ **Indications** Adrenal insufficiency, orthostatic hypotension in diabetics

⊙ **Common drug examples**

- **Fludocortisone acetate** Florinef

⊙ **Adverse reactions** Salt and water retention, hypertension, cardiac hypertrophy, edema, heart failure, bruising, diaphoresis, urticaria, allergic rash, hypokalemia

[Note: All adrenocorticoid drugs have both glucocorticoid and mineralocorticoid properties to some extent]

alpha-adrenergic blockers

Lower blood pressure by dilating peripheral blood vessels, reducing peripheral resistance

⊙ **Indications** Raynaud's disease, acrocyanosis, frostbite, phlebitis, diabetic gangrene, hypertension, benign prostatic hyperplasia

⊙ **Common drug examples:**

- **Doxazosin mesylate** Cardura
- **Prozosin hydrochloride** Minipress
- **Tamsulosin hydrochloride** Flomax
- **Terazosin hydrochloride** Hytrin

⊙ **Adverse reactions** Orthostatic hypotension, headache, palpitations, fatigue, nausea, weakness, dizziness, fainting

aminoglycosides

⊙ **Indications:**

- Treat infections resistant to penicillin, septicemia, urinary tract infections, infections of skin, soft tissue and bone, gram-negative bacillary meningitis
- Used in combination with other antibiotics to treat staphylococcal infections, endocarditis, tuberculosis, pelvic inflammatory disease

⊙ **Common drug examples:**

- **Amikacin sulfate** Amikin
- **Gentamicin sulfate** Cidonycin, Gentsol
- **Neomycin sulfate** Mycifradin

⊙ **Adverse reactions** Systemic ototoxicity and nephrotoxicity, skeletal weakness and respiratory distress; oral meds can cause nausea, vomiting, diarrhea; local injections can cause phlebitis and abscess

androgens

Testosterone used to promote maturation of male sex organs and development of secondary sex characteristics; promotes retention of calcium, nitrogen, phosphorus, sodium, and potassium; enhances anabolism

⊙ **Indications** Androgen deficiency resulting from testicular failure or deficiency of pituitary origin, palliative for metastatic breast cancer, postpartum breast engorgement, hereditary angioedema, endometriosis, fibrocystic breast disease

⊙ **Common drug examples:**

- **Danazol** Cyclomen, Danocrine
- **Fluoxymesterone** Halotestin
- **Testosterone** Testopel pellets

⊙ **Adverse reactions:**

- **Extensions of hormonal action**
 - **Males:** Frequent and prolonged erections, bladder irritability, gynecomastia
 - **Females:** Clitoral enlargement, deepening of the voice, facial or body hair growth, unusual hair loss, irregular or absent menses
- **Metabolic reactions** Fluid and electrolyte retention, hypercalcemia, decreased blood glucose level, increased serum cholesterol, hepatic dysfunction

⊙ **Contraindicated** Men with breast or prostatic cancer or symptomatic prostate hypertrophy, patients with severe cardiac, renal or hepatic disease or with undiagnosed genital bleeding

angiotensin-converting enzyme inhibitors

⊙ **Indications** Treat high blood pressure and heart failure

⊙ **Common drug examples:**

- **Benazepril hydrochloride** Lotensin
- **Captopril** Capoten
- **Enalapril maleate** Vasotec
- **Fosinopril sodium** Monopril
- **Lisinopril** Prinivil, Zestril

⊙ **Adverse reactions** Persistent dry cough, skin rash, loss of taste, weakness, headaches, palpitations, fatigue, proteinuria, hyperkalemia

angiotensin II receptor antagonists

Vasodilates arterioles by blocking the effects of angiotensin II, enhance renal clearance of sodium and water

⊙ **Indications** Treatment of high blood pressure

⊙ **Common drug examples:**

- **Candesartan cilexetil** Atacand
- **Eprosartan mesylate** Teveten
- **Irbesartan** Avapro
- **Losartan potassium** Cozaar
- **Telmisartan** Micardis
- **Valsartan** Diovan

⊙ **Adverse reactions** Dizziness, anxiety, confusion, cough, upper respiratory infections, myalgia, insomnia, hypotension, visual changes, GI/GU effects

anticholinergics

⊙ **Indications:**

- Spastic conditions including Parkinson's disease, muscle dystonia, muscle rigidity and extra-pyramidal disorders

- Prevent nausea and vomiting from motion sickness, adjunctive treatment for peptic ulcers and other GI disorders, bronchospasms, and GU tract disorders
- Treat poisoning from certain plants and pesticides
- Use preoperatively to decrease secretions and block cardiac reflexes
- ⊙ **Common drug examples:**
 - **Antiparkinsonians:**
 - *Benztropine mesylate*: cogentin
 - **Belladonna alkaloids:**
 - *Scopolamine hydrobromide*: IsoptoHyoscine, Scopace
 - **Synthetic quaternary anticholinergics:**
 - *Glycopyrrrolate*: Robinul
 - **Tertiary synthetic and semisynthetic derivatives:**
 - *Dicyclomine hydrochloride*: Antispas, A-spas, Dibent, Dilomine, Lomine, Ortyl
- ⊙ **Adverse reactions** Dry mouth, decreased sweating, headache, dilated pupils, blurred vision, dry skin, urinary hesitancy and urine retention, constipation, palpitations and tachycardia; other peripheral effects include dry mucous membranes, dysphasia, stupor, seizures, hyperthermia, hypertension and increased respiration
- **Toxic doses** May cause disorientation, confusion, hallucinations, delusions, anxiety, agitation and restlessness

anticoagulants

- ⊙ **Indications** Prevent clot formation in patients with DVTs and pulmonary embolism, provide anticoagulation during hemodialysis, prevention of postoperative clot formation after surgery, decrease risk of strokes, decrease risk of MI in patients with atherosclerosis
- ⊙ **Common drug examples:**
 - **Danaparoid** Orgaran
 - **Delteparin** Fragmin
 - **Enoxaparin** Lovenox
 - **Heparin** Heparin Lock Flush, Hep-lock
 - **Tinzaparin** Innohep
- ⊙ **Adverse reactions** Insomnia, headache, dizziness, confusion, peripheral edema, nausea, constipation, pain, fever, vomiting, joint pain, rash

antihistamines

- ⊙ **Indications** Allergies, pruritis, vertigo, nausea and vomiting, sedation, suppression of cough, dyskinesia
- ⊙ **Common drug examples:**
 - **Allergies:**
 - *Azelastine hydrochloride*: Astelin, Optivar
 - *Chlopheniramine maleate*: Aller-Chlor, Chlor-Trimeton, Chlor-Tripolon
 - *Clemastine fumarate*: Tavist
 - *Diphenhydramine hydrochloride*: Allergy DM, Benadryl, Diphen, Dormin, Midol PM, Nytol, Sominex, Twilite
 - *Promethazine hydrochloride*: Anergan 50, Phenergan
 - **Pruritis:**
 - *Cyproheptadine hydrochloride*: Periactin
 - *Hydroxyzine hydrochloride*: Anxanil, Atarax, Multipax, Quiess, Vistacon
 - **Vertigo, nausea, vomiting:**
 - *Cyclizine hydrochloride*: Marezine
 - *Cyclizine lactate*: Marezine, Marzine
 - *Dimenhydrinate*: Dimetab, Hydrate, Triptone
 - *Meclizine hydrochloride*: Antivert, Antrizine, Bonine, Vergon
 - *Promethazine hydrochloride*: Anergan, Phenergan
 - **Sedation:**
 - *Diphenhydramine*: Diphenhydramine syrup
 - **Cough suppression:**
 - Diphenhydramine syrup
 - **Dyskinesia:**
 - Diphenhydramine
- ⊙ **Adverse reactions** Drowsiness and impaired motor function; anticholinergic action will cause dry mouth and throat, blurred vision and constipation
- **Toxic effects** Sedation, reduced mental alertness, apnea, cardiovascular collapse, hallucinations, tremors, seizures, dry mouth, flushed skin, and fixed, dilated pupils; (reverses when drug is withdrawn)

anxiolytic skeletal muscle relaxant

- ⊙ **Indications** Anxiety, muscle spasm, tetanus, acute alcohol withdrawal, adjunct for epilepsy
- ⊙ **Common drug examples:**
 - **Diazepam** Apo-Diazepam, Valium
- ⊙ **Adverse reactions** Drowsiness, slurred speech, tremor, fatigue, ataxia, headache, insomnia, hypotension, bradycardia, nausea, constipation, joint pain, physical or psychological dependence

barbiturates

- ⊙ **Indications** Seizure disorders (tonic-clonic and partial seizures), sedation, hypnosis, preanesthesia sedation, psychiatric use

⊙ **Common drug examples:**

- **Amobarbital** Amytal
- **Phenobarbital** Bellatal, Solfoton
- **Primidone** Mysoline
- **Secobarbital sodium** Seconal

⊙ **Adverse reactions** Drowsiness, lethargy, vertigo, headaches and CNS depression, hypersensitivity can occur (rash, fever)

- **After hypnosis** hangover effect, impaired judgment, mood distortion, rebound insomnia
- **Geriatric patients** Confusion
- **Pediatric patients** Hyperactivity

benzodiazepines

Enhance/facilitate actions of the gamma-aminobutyric acid (GABA)

⊙ **Indications** Seizure disorders, anticonvulsants, anxiety, tension and insomnia, surgical adjuncts for conscious sedation or amnesia, skeletal muscle spasms or tremors, delirium, schizophrenia as an adjunct, nausea and vomiting induced by chemotherapy, neonatal opiate withdrawal

⊙ **Common drug examples:**

- **Alprazolam** Alprazolam, Xanax
- **Chlordiazepoxide** Libritab
- **Clonazepam** Klonopin, Rivotril
- **Clorazepate dipotassium** Catapres, Dixarit
- **Diazepam** Valium, Zetran
- **Estazolam** ProSom
- **Flurazepam** Apo-Flurazepam, Dalmane
- **Lorazepam** Apo-Lorazepam, Ativan
- **Midazolam** Versed
- **Oxazepam** Apo-Oxazepam, Serax
- **Temazepam** Restoril
- **Triazolam** Halcion

⊙ **Adverse reactions** Drowsiness and impaired motor function; constipation, diarrhea, vomiting, changes in appetite, urinary alterations, nightmares, hallucinations, insomnia

- **Toxic effects** Visual disturbances, short-term memory loss, vertigo, confusion, severe depression, shakiness, slurred speech, staggering, bradycardia, difficulty breathing

beta blockers

Reduce the workload of the heart by blocking the sympathetic conductance at the beta receptors on the SA node and myocardial cells, thus decreasing the force of contraction and causing a reduction in heart rate

⊙ **Indications** Hypertension, angina, arrhythmias, glaucoma, myocardial infarction, migraine prophylaxis

⊙ **Common drug examples:**

- **Beta 1 Blockers:**
 - *Acebutolol*: Sectral
 - *Atenolol*: Tenormin
 - *Betaxolol hydrochloride*: Betoptic, Kerlone
 - *Bisoprolol fumarate*: Zebeta
 - *Esmolol*: Brevibloc
 - *Metoprolol tartrate*: Lopressor
- **Beta 1 & 2 Blockers:**
 - *Carteolol*: Cartrol, Ocupress
 - *Carvedilol*: Coreg
 - *Labetalol hydrochloride*: Normodyne, Trandate
 - *Levobunolol hydrochloride*: AKBeta, Betagen
 - *Metipranolol hydrochloride*: Opti Pranolol
 - *Nadolol*: Corgard
 - *Pindolol*: Visken
 - *Propranolol*: Inderal
 - *Sotalol*: Betapace
 - *Timolol maleate*: Blocarden, Timoptic

⊙ **Adverse reactions** Insomnia, nausea, fatigue, slow pulse, weakness, increased cholesterol and blood glucose levels, bradycardia, depression, hallucinations, sexual dysfunctions, skin hyperpigmentation

- **Toxic effects** Severe hypotension, bradycardia, heart failure, bronchospasms

bile acid sequestrants

⊙ **Indications** Lowering cholesterol

⊙ **Common drug examples:**

- **Cholestyramine** Locholest, Prevalite, Questran
- **Colesevelam** Welchol
- **Colestipol** Colestid

⊙ **Adverse reactions** Headache, anxiety, vertigo, dizziness, insomnia, fatigue, syncope, tinnitus, constipation, nausea, vomiting, anemia, muscle and joint pain

calcium channel blockers

Relaxes smooth muscle to provide vasodilation and affects cardiac muscle to reduce HR and SV

- ⊙ **Indications** Angina, arrhythmias, hypertension, migraine headaches, peripheral vascular disorders, subarachnoid hemorrhage, esophageal spasm (adjunctive therapy)
- ⊙ **Common drug examples:**
 - **Amlodipine besylate** Norvasc
 - **Bepidil hydrochloride** Vascor
 - **Diltiazem hydrochloride** Cardizem, Dilacor, Tiazac
 - **Felodipine** Plendil
 - **Isradipine** DynaCirc
 - **Nicardipine** Cardene
 - **Nifedipine** Procardial
 - **Nimodipine** Nimotop
 - **Nisoldipine** Sular
 - **Verapamil hydrochloride** Calan, Isoptin, Verelan
- ⊙ **Adverse reactions** Bradycardia, hypotension, fluid retention, palpitations, headaches from vasodilatation, flushes, rash, dizziness
 - Verapamil can cause constipation
 - Nifedipine can cause hypotension, reflex tachycardia, peripheral edema, flushing, light-headedness and headache
 - Diltiazem can cause anorexia, nausea, heart block, bradycardia, heart failure and peripheral edema

cephalosporin

Antibiotics that inhibit bacterial cell wall synthesis, causing bacterial cell death

- ⊙ **Indications** Serious infections of the lungs, skin, soft tissue, bones, joints, urinary tract, blood (septicemia), abdomen and heart (endocarditis), second and third generation drugs can treat CNS infections (meningitis), Lyme disease
- ⊙ **Common drug examples:**
 - **First Generation:**
 - **Cefadroxil:** Duricef
 - **Cefazolin sodium:** Ancef, Defzol
 - **Cephalexin monohydrate:** Biocef, Keflex, Novo-Lexin
 - **Cephadrine:** Keftab
 - **Second Generation:**
 - **Cefaclor:** Ceclor
 - **Cefamandole nafate:** Mandol
 - **Cefotetan disodium:** Cefotan
 - **Cefprozil:** Cefzil
 - **Cefuroxime axetil:** Ceftin
 - **Cefuroxime sodium:** Kefurox, Zinacef
 - **Third Generation:**
 - **Cefdinir:** Omnicef
 - **Cefditoren pivoxil:** Spectracef
 - **Cefixime:** Suprax
 - **Cefoperazone sodium:** Cefobid
 - **Cefotaxime sodium:** Claforan
 - **Cefpodoxime proxetil:** Vantin
 - **Ceftazidime:** Ceptaz, Fortaz, Taxicef, Taxidime
 - **Ceftizoxime sodium:** Cefizox
 - **Ceftriaxone sodium:** Rocephin
 - **Fourth Generation:**
 - **Cefepime hydrochloride:** Maxipime
- ⊙ **Adverse reactions** Mild rash, fever, fatal anaphylaxis (hypersensitivity); thrombocytopenia, transient neutropenia, reversible leucopenia; other effects include nausea, vomiting, diarrhea, abdominal pain, glossitis, dyspepsia; local venous pain and irritation are common at injection site

diuretics

Loop Increase the excretion of sodium and water and control high blood pressure and fluid retention

- ⊙ **Indications** Edema associated with heart failure, hypertension, renal impairment, hypertensive crisis
 - ⊙ **Common drug examples:**
 - **Bumetanide** Bumex
 - **Ethacrynic acid** Endecrin
 - **Furosemide** Lasix
 - **Torsemide** Demadex
 - ⊙ **Adverse reactions** Metabolic and electrolyte disturbances, hypochloremic alkalosis, hyperglycemia, hyperuricemia, hypomagnesemia, may cause hearing loss and tinnitus
- Potassium-sparing Less potent than the other types, protects against potassium loss*
- ⊙ **Indications** Edema associated with hepatic cirrhosis, nephritic syndrome, and heart failure, hypertension, primary hyperaldosteronism
 - ⊙ **Common drug examples:**
 - **Amiloride hydrochloride** Midamor
 - **Spironolactone** Aldactone
 - **Triamterene** Dyrenium
 - ⊙ **Adverse reactions** Hyperkalemia leading to arrhythmias, nausea, vomiting, headaches, weakness, fatigue, bowel disturbances, cough and dyspnea

Thiazide

- ⊙ **Indications** Edema caused by heart failure and nephritic syndromes, edema caused by pregnancy, hypertension, diabetes insipidus
- ⊙ **Common drug examples:**
 - **Bendroflumethiazide** Naturetin
 - **Chlorothiazide** Diuril
 - **Chlorthalidone** Hygroton
 - **Hydrochlorothiazide** Esidrix, HydroDiuril, Microzide, Oretic

- **Hydroflumethiazide** Diucardin
- **Indapamide** Lozol
- **Methychothiazide** Aquatensen, Enduron
- **Metolazone** Mykrox, Zaroxolyn
- **Trichlormethiazide** Diurese, Metahydrin, Naqua
- ⊙ **Adverse reactions** Electrolyte and metabolic disturbances, hypochloremic alkalosis, hypomagnesemia, hyponatremia, hypercalcemia, hyperuricemia, elevated cholesterol levels, hyperglycemia, lethargy (overdose can progress to coma)

estrogens

- ⊙ **Indications** Menopause, carcinoma of the prostate, cardiovascular risk prevention, prophylaxis of postmenopausal osteoporosis, contraception, some drugs are used in treatment of breast cancer—must be carefully selected patients and drugs
- ⊙ **Common drug examples:**
 - **Dienestrol** Orthodienestrol vaginal cream
 - **Esterified estrogen** Estrab, Menest
 - **Estradiol** Alora, Climara, Esclim, Estrace, Estraderm, Estring, Fem Patch, Vivelle
 - **Estradiol cypionate** Depo-Estradiol Cypionate, DepGynogen, DepoGen
 - **Estradiol valerate** Delestrogen, Gynogen LA 20, Valergen 20, Valergen 40
 - **Ethinyl estradiol** Estinyl
 - **Estropipate** Ogen, Ortho-Est
- ⊙ **Adverse reactions** Menstrual bleeding, abdominal cramps, swollen feet or ankles, bloated sensation, breast swelling and tenderness, weight gain, nausea, loss of appetite, headaches, photosensitivity and loss of libido; long-term use can cause hypertension, thromboembolic disease

fluoroquinolones

Antibacterial agent used against aerobic gram-positive and gram-negative organisms

- ⊙ **Indications** Bone and joint infections, bacterial bronchitis, endocervical and urethral chlamydia, bacterial gastroenteritis, endocervical and urethral gonorrhea, intra-abdominal infections, empiric therapy for febrile neutropenia, pelvic inflammatory disease, bacterial pneumonia, bacterial prostatitis, acute sinusitis, skin and soft tissue infections, typhoid fever, bacterial urinary tract infections, chancroid, meningococcal carriers, bacterial septicemia, prophylaxis in prevention of bacterial urinary tract infections
- ⊙ **Common drug examples:**
 - **Ciprofloxacin** Ciloxan
 - **Esylate/alatrofloxacin mesylate** Trovan IV
 - **Gatifloxacin** Tequin
 - **Levofloxacin** Quixin
 - **Lomefloxacin hydrochloride** Maxaquin
 - **Moxifloxacin hydrochloride** Avelox
 - **Norfloxacin** Chibroxin
 - **Ofloxacin** Floxin, Ocuflax
 - **Sparfloxacin** Zagam
 - **Trovafloxacin** Trovan
- ⊙ **Adverse reactions** Rarely seen; acute stimulation of the CNS causes acute psychosis, agitation, hallucinations and tremors; hepatotoxicity, tendonitis or tendon rupture; other effects include dizziness, headache, nervousness, drowsiness, insomnia, GI reactions and photosensitivity

histamine-receptor antagonists

- ⊙ **Indications** Duodenal ulcer, gastric ulcer, hypersecretory states, acid reflux, esophagitis, stress ulcer prophylaxis
- ⊙ **Common drug examples:**
 - **Cimetidine** Tagemet
 - **Famotidine** Pepcid, Pepcid AC
 - **Nizatidine** Apo-Nizatidine, Axid
 - **Ranitidine** Zantac
 - **Rantidine bismuth citrate**
- ⊙ **Adverse reactions** Mild transient diarrhea, neutropenia, dizziness, fatigue, arrhythmias, gynecomastia

HMG-CoA reductase inhibitors

- ⊙ **Indications** Hyper cholesterol, mixed dyslipidemia, secondary prevention of cardiovascular events (except atorvastatin)
- ⊙ **Common drug examples:**
 - **Atorvastatin** Lipitor
 - **Fluvastatin sodium** Lescol, Lescol XL
 - **Lovastatin** Mevacor
 - **Pravastatin sodium** Pravachol
 - **Simvastatin** Zocor
- ⊙ **Adverse reactions** Photosensitivity, hepatotoxicity, GI complaints, myopathy (usually muscle aches and weakness), insomnia

leukotriene receptor blockers

- ⊙ **Indications** Asthma
- ⊙ **Common drug examples:**
 - **Montelukast Sodium** Singulair
 - **Zileuton** Zyflo
 - **Zafirlukast** Accolate

- ⊙ **Adverse reactions** Headache, dizziness, fatigue, fever, dyspepsia, asthenia, abdominal pain, cough, influenza, pain

nitrates

- ⊙ **Indications** Angina pectoris, acute myocardial infarction, hypertensive emergencies, heart failure and pulmonary edema associated with MI
- ⊙ **Common drug examples:**
 - **Isosorbide dinitrate** Apo-ISDN, Coronex, Isoril, Novosorbide, Sorbitrate
 - **Isosorbide mononitrate** Imdur, ISMO, Isotrate ER, Monoket
 - **Nitroglycerin** Nitro-Bid, Nitrogly, Nitrong, Nitrostat, Nitrolingual
- ⊙ **Adverse reactions** Headaches, orthostatic hypotension, dizziness, weakness and transient flushing, nausea, vomiting, restlessness, pallor, cold sweats, tachycardia, syncope or CV collapse can occur

non-steroidal anti-inflammatory (NSAIDs)

Analgesic, anti-inflammatory, antipyretic

- ⊙ **Indications** Pain, inflammation, and fever; rheumatoid arthritis, juvenile arthritis and osteoarthritis; low-intensity headaches, arthralgia, myalgia, neuralgia and mild to moderate pain from dental or surgical procedures or dysmenorrhea
- ⊙ **Common drug examples: (OTC and prescription)**
 - **Acetylsalicylic Acid (ASA)** Aspirin
 - **Celecoxib** Celebrex
 - **Diclofenac sodium** Voltaren
 - **Diflunisal** Dolobid
 - **Etodolac** Lodine
 - **Fenoprofen calcium** Nalfon
 - **Ibuprofen** Advil, Medipren, Motrin, Nuprin, Rufen, Trendar
 - **Indomethacin** Indocid, Novomethacin
 - **Ketoprofen** Oradis, Oruvail
 - **Ketorolac tromethamine** Toradol
 - **Nabumetone** Relafen
 - **Naproxen** Naprosyn
 - **Naproxen sodium** Anaprox, Aleve, Naprelan
 - **Oxaprozin** Daypro
 - **Piroxicam** Feldene
 - **Sulindac** Clinoril
 - **Tolmetin sodium** Tolectin
- ⊙ **Adverse reactions** Abdominal pain, bleeding, anorexia, ulcers, liver toxicity, dyspepsia, heartburn (minimized if taken with meals); flank pain may indicate nephrotoxicity; drowsiness, headache, dizziness, confusion, tinnitus, vertigo, depression, bladder infections, blood in urine and kidney necrosis

nucleoside reverse transcriptase inhibitors

- ⊙ **Indications** Used in combination with other drugs to treat HIV infections and AIDS; prevention of maternal/fetal HIV transmission, prevention of HIV infection after an occupational exposure
- ⊙ **Common drug examples:**
 - **Abacavir sulfate** Ziagen
 - **Didanosine** Videx
 - **Lamivudine** Combivir
 - **Stavudine** Zerit
 - **Zalcitabine** Hivid
 - **Zidovudine** Retrovir
- ⊙ **Adverse reactions** Difficult to distinguish between disease-related side effects and drug-related side effects; anemia, leucopenia, neutropenia, thrombocytopenia
- **Toxic effects** Rare adverse effects that require medical attention: myopathy, neurotoxicity and hepatotoxicity; not requiring medical attention: headache, severe insomnia, myalgia, nausea or hyperpigmentation of nails

opioids (previously called narcotics)

Often used in combination with other medications, particularly acetaminophen

- ⊙ **Indications** Analgesic used for moderate to severe pain associated with acute and chronic disorders including MI, postoperative pain or terminal cancer; pulmonary edema, preoperative sedation, anesthesia, cough suppression, diarrhea
- ⊙ **Common drug examples:**
 - **Codeine phosphate; codeine sulfate**
 - **Diphenoxylate hydrochloride** Lofene, Lomotil
 - **Fentanyl citrate** Sublimaze
 - **Fentanyl transdermal system** Duragesic
 - **Meperidine hydrochloride** Demerol
 - **Methadone hydrochloride** Dolphine, Mehadose
 - **Morphine sulfate** Epimorph, Kadian, Statex
 - **Oxycodone hydrochloride** Endocodone, Percolone
- ⊙ **Adverse reactions** Respiratory depression, circulatory depression, respiratory arrest, cardiac arrest, dizziness, visual disturbances, mental clouding, sedation, coma, euphoria, weakness, agitation, restlessness, nervousness, seizures, nausea, vomiting, constipation; high potential for addiction

opioid mixed agonist-antagonist

Drugs have both agonistic and antagonistic properties; usually potent analgesics but less addictive than pure opioids

- ⊙ **Indications** [See Opioids]
- ⊙ **Common drug examples:**
 - **Buprenorphine hydrochloride** Buprenex
 - **Butorphanol tartrate** Stadol
 - **Nalbuphine hydrochloride** Nubain
 - **Pentazocine hydrochloride** Talwin
- ⊙ **Adverse reactions** Respiratory depression, apnea, shock and cardiopulmonary arrest; sedation, dizziness, hallucinations, disorientation, agitation, euphoria, dysphoria; insomnia; headache; miosis, tachycardia, palpitations, chest wall rigidity, syncope and edema; nausea, vomiting and constipation; dry mouth; anorexia and spasms of the colon; urinary retention or hesitancy; decreased libido; rash, flushing; physical and psychological dependence can occur

penicillin

Family of effective antibiotics with low toxicity

- ⊙ **Indications:**
 - **Natural penicillin** Infections like streptococcal pneumonia, enterococcal and nonenterococcal group D endocarditis, diphtheria, anthrax, meningitis, tetanus, botulism, actinomycosis, syphilis, and relapsing fever, Lyme disease; prophylaxis against pneumococcal infections, rheumatic fever, bacterial endocarditis
 - **Aminopenicillins** Septicemia; gynecologic infections; respiratory, GU and GI tract infections, soft tissue, bone and joint infections
 - **Extended-spectrum penicillins:** Hard to treat gram-negative infections; given in combination with aminoglycosides
 - **Penicillinase-resistant penicillins:** Susceptible penicillinase producing staphylococci; much the same as for aminopenicillins
- ⊙ **Common drug examples:**
 - **Natural penicillin:**
 - **Penicillin G benzathine:** Bicillin L-A, Permapen
 - **Penicillin G potassium:** Pfizerpen
 - **Penicillin G procaine:** Bicillin C-R, Wycillin
 - **Penicillin V potassium:** Apo-Pen, Veetids
 - **Aminopenicillins:**
 - **Amoxicillin trihydrate with clavulanate potassium:** Augmentin, Clavulin
 - **Ampicillin:** Apo-Ampi Novo-Ampicillin, Omnipen, Penbritin
 - **Ampicillin trihydrate:** Principen, Totacillin
 - **Penicillinase-resistant penicillins:**
 - **Dicloxacillin sodium:** Dycil, Dynapen, Pathocil
 - **Nafcillin sodium:** Nafcil, Nailpen, Unipen
 - **Oxacillin sodium:** Bactocil
 - **Mezlocillin sodium:** Mexlin
 - **Piperacillin sodium:** Pipracil
 - **Piperacillin sodium with tazobactam sodium:** Zosyn
 - **Ticarcillin disodium:** Ticar
 - **Ticarcillin with clavulanate potassium:** Timentin
- ⊙ **Adverse reactions** Hypersensitivity reactions, hematological reactions, transient neutropenia, leucopenia, thrombocytopenia; bleeding can occur with high-dose extended-spectrum penicillins

phenothizines

- ⊙ **Indications** Psychoses involving hallucinations, agitation, manic phase of bipolar psychoses; nausea and vomiting induced by CNS dysfunctions; anxiety; severe behavioral problems, abdominal pain associated with porphyria, delirium, neurogenic pain
- ⊙ **Common drug examples:**
 - **Aliphatic derivatives:**
 - **Chlorpromazine hydrochloride:** Chlorpromanyl—20, Largac-til, Thorazine
 - **Promethazine hydrochloride:** Anergan 50, Phenergan
 - **Piperazine derivatives:**
 - **Fluphenazine hydrochloride:** Permitil, Prolixin
 - **Perphenazine:** Apo-Perphenazine, Trilafon
 - **Prochlorperazine:** Compazine, Stermetil
 - **Trifluoperazine hydrochloride:** Apo-Trifluoperazine, Stelazine
 - **Piperidine derivatives:**
 - **Mesoridazine besylate:** Serentil
 - **Thioridazine:** Mellaril-S
 - **Thiothixene hydrochloride:** Navane
- ⊙ **Adverse reactions** Some medications may cause extra-pyramidal symptoms; in rare cases can cause neuroleptic malignant syndrome; other reactions include sedative and anticholinergic effects, orthostatic hypotension, reflex tachycardia, fainting, dizziness, arrhythmias, anorexia, nausea, vomiting, local gastric irritation, endocrine effects, hematological disorders, ocular changes

progestins

- ⊙ **Indications** Hormonal imbalance in women, endometriosis, carcinoma, contraception

Common drug examples:

- **Medroxyprogesterone acetate** Amen, Curretab, Cycrin, Provera
- **Megestrol acetate** Megace
- **Norethindrone** Micronor, Nor-Q.D.
- **Norethindrone acetate** Aygestin, Norlutate
- **Norgestrel** Ovrette
- **Progesterone** Crinone

- ⊙ **Adverse reactions** Change in menstrual bleeding pattern, breast tenderness and secretion, weight changes, increases in body temperature, edema, nausea, acne, somnolence, insomnia, hirsutism, hair loss, depression, cholestatic jaundice and allergic reactions; flushing, increased sugar levels, increase in BP, decreased sexual desire, headache

protease inhibitors

Antiviral medication used with HIV patients

- ⊙ **Indications** HIV infection and AIDS

Common drug examples:

- **Amprenavir** Agenerase
- **Indinavir sulfate** Crixivan
- **Lopinavir and ritonavir** Kaletra
- **Nelfinavir mesylate** Viracept
- **Ritonavir** Norvir
- **Saquinavir** Fortovase
- **Saquinavir mesylate** Invirase

- ⊙ **Adverse reactions** Kidney stones, pancreatitis, diabetes or hyperglycemia, ketoacidosis and paresthesia all require medical attention; less problematic are symptoms of generalized weakness, GI disturbances, headaches, insomnia, taste perversion, dizziness, somnolence

selective serotonin reuptake inhibitors

Enhance serotonergic transmission through blocked reuptake at the synapse

- ⊙ **Indications** Depression, panic and eating disorders, obsessive compulsion, premenstrual dysphoria, posttraumatic stress and bipolar disorders, alcohol dependence, premature ejaculation, diabetic neuropathy

Common drug examples:

- **Citalopram hydrobromide** Celexa
- **Fluoxetine** Prozac, Sarafem
- **Fluvoxamine maleate** Luvox
- **Paroxetine hydrochloride** Paxil
- **Sertraline hydrochloride** Zoloft

- ⊙ **Adverse reactions** GI complaints, headaches, dizziness, somnolence, sexual dysfunction, tremors; less common reactions include breast tenderness or enlargement, extra-pyramidal effects, dystonia, fever, palpitations, weight gain or loss, rash, hives, itching

skeletal muscle relaxant I

Polysynaptic inhibitors (inhibit interneuron transmission in the spinal cord)

- ⊙ **Indications** Muscle spasms caused by acute injuries, supportive therapy for tetanus

Common drug examples:

- **Carisoprodol** Soma
- **Chlorzoxazone** Paraflex, Parafon Forte
- **Cyclobenzaprine hydrochloride** Flexeril
- **Methocarbamol** Carbacot, Robaxin, Skelex
- **Orphenadrine citrate** Norflex

- ⊙ **Adverse reactions** Drowsiness, vertigo, tremor, headaches, light-headedness, nausea, vomiting, confusion

skeletal muscle relaxant II

Indirect and direct skeletal muscle relaxants

- ⊙ **Indications** Spasticity caused by an upper motor neuron lesion like MS

Common drug examples:

- **Baclofen** Lioresal
- **Diazepam** Valium
- **Dantrolene sodium** Dantrium

- ⊙ **Adverse reactions** Drowsiness, dizziness, weakness, fatigue, hypotension, paresthesias, confusion, dysarthria, constipation, vomiting, liver dysfunction

sulfonamides

First drugs to treat systemic, bacterial infections

Indications:

- **Bacterial infections** Effective with staphylococci, streptococci, clostridium tetani, urinary tract infections, nocardiosis, otitis media
- **Parasitic infections** Inflammation, pneumonic plague

Common drug examples:

- **Co-trimoxazole** Apo-Sulfatrim, Bactrim, Cotrim, Septra
- **Sulfasalazine** Azulfidine

- ⊙ **Adverse reactions** Rash, fever, pruritus, erythema, photosensitivity, joint pain, bronchospasm; hematologic, renal and GI reactions all can occur

sulfonylureas

Lower blood glucose levels by stimulating insulin release from the pancreas

- ⊙ **Indications** Type 2 diabetes mellitus, neurogenic diabetes insipidus

Common drug examples:

- **First Generation:**
 - Chlorpropamide: Diabinese, Novo-propamide
 - Tolazamide: Tolinase
 - Tolbutamide: Orinase
- **Second Generation:**
 - Glimepiride: Amaryl
 - Glipizide: Glucotrol
 - Glyburide: DiaBeta, Glynase Pres Tab, Micronase

- ⊙ **Adverse reactions** Headache, nausea, vomiting, anorexia, heartburn, weakness and paresthesia

- **Toxic effects** Anxiety, chills, cold sweats, confusion, cool pale skin, difficulty concentrating, drowsiness, excessive hunger, nervousness, rapid heartbeat, weakness, unusual fatigue

tetracycline

Antibiotic

- ⊙ **Indications** Bacterial, antiprotozoal, rickettsial and fungal infections; sclerosing agent for pleural or pericardial effusion, adjunct therapy for H. pylori and other GI infections, Lyme disease

Common drug examples:

- **Doxycycline hyclate** Periostat, Vibramycin
- **Minocycline hydrochloride** Dynacin, Nimocin, Vectrin
- **Tetracycline hydrochloride** Achromycin, Panmycin, Tetralen

- ⊙ **Adverse reactions** Anorexia, flatulence, nausea, vomiting, stool disturbances, epigastric burning, abdominal discomfort, rash

thrombolytic enzymes

Developed to reduce a blood clot and prevent permanent ischemic damage

- ⊙ **Indications** Thrombosis, thromboembolism

Common drug examples:

- **Alteplase** Activase, Cathflo Activase
- **Anistreplase, reteplase** Eminase
- **Streptokinase** Streptase
- **Tenecteplase** TNKase
- **Urokinase** Abbokinase

- ⊙ **Adverse reactions** Cerebral hemorrhage, fever, hypotension, arrhythmias, edema, nausea, vomiting, arthralgia, headache

tricyclic antidepressants

Enhance adrenergic neurotransmitter transmission through blocked reuptake at the synapse

- ⊙ **Indications** Depression, obsessive compulsive disorder, enuresis, severe chronic pain, phobic disorders, bulimia, short-term treatment of duodenal or gastric ulcers

Common drug examples:

- **Amitriptyline hydrochloride** Elavil, Levate, Novotriptyn
- **Clomipramine hydrochloride** Anafranil
- **Desipramine hydrochloride** Norpramin
- **Doxepin hydrochloride** Sinequan, Triadapin
- **Imipramine hydrochloride** Apo-Imipramine, Impril, Novopramine
- **Imipramine pamoate** Tofranil-AM
- **Nortriptyline hydrochloride** Aventyl HCL, Pamelor
- **Trimipramine maleate** Surmontil

- ⊙ **Adverse reactions** Sedation, anticholinergic effects, orthostatic hypotension; specific drugs may cause seizures

vitamin K inhibitors

- ⊙ **Indications** Pulmonary emboli, DVT, MI, atrial arrhythmias

Common drug examples:

- **Warfin** Coumadin

- ⊙ **Adverse reactions** Fever, anorexia, nausea, vomiting, cramps, diarrhea, mouth ulcerations, hemorrhage, jaundice

U.S. \$5.95 CAN. \$8.95

NOTE TO STUDENTS Due to its condensed format, please use this QuickStudy® as a guide, but not as a replacement for assigned classwork.

DISCLAIMER: This guide cannot cover every possible adverse reaction or toxic effect, and is intended for informational purposes only; it is not intended for the diagnosis, treatment or cure of any medical condition or illness, and should not be used as a substitute for professional medical care. BarCharts, Inc., its writers and editors are not responsible or liable for the use or misuse of the information contained in this guide.

free downloads & hundreds of titles at quickstudy.com

ISBN-13: 978-142320749-8
ISBN-10: 142320749-1

9 781423 120749 8

5 9 5 9 5

Authors: Becky Rodda, PT, MHS, OCS
Suzanne L. Tinsley, PhD, PT

Customer Hotline # 1.800.230.9522

All rights reserved. No part of this publication may be reproduced or transmitted in any form, or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission from the publisher.
©2006 BarCharts, Inc. 0608

6 54614 100749 0